# Introducing two new SUPER GUARD formulas designed to fit your individual calf management program needs!

SUPER GUARD - Regular formulation containing the maximum allowable dosage of Neo/Oxy. SUPER GUARD PLUS - Regular formulation with the added benefits of Direct Fed Microbials, Kaolin and high levels of Vitamins A, D3 and E.

> Convenient feeding directions provide the maximum allowable amount of 10 mg/lb of body weight per day of Neomycin Sulfate and Oxytetracyline Hydrochloride.



For treatment of bacterial enteritis (scours) caused by *E. coli* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *E. coli* susceptible to neomycin.

For treatment of bacterial pneumonia (shipping fever complex) caused by Pasteurella

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate .......
Oxytetracycline Hydrochloride

Providing the maximum allowable amount of 10 mg/lb of body weight per day, when fed as directed.

#### **GUARANTEED ANALYSIS**

Crude Protein, minimum	6.0%
Crude Fat, minimum	6.0%
Crude Fiber, maximum	0.15%
Calcium, minimum	0.2%
Calcium, maximum	0.5%
Phosphorus, minimum	0.4%

#### INGREDIENTS

Dried Whey, Maltodextrin, Dextrose, Animal Fat (preserved with BHA/BHT) and Natural and Artificial Flavors.

Keep tightly closed. Store in cool, dry place.

#### WARNING

Residue Warning: Withdrawal 5 days before slaughter. Use of more than one product containing neomycin or failure to follow withdrawal times may result is illegal drug residue. A withdrawal period has not been established for the use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for the use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

For Animal Use Only. Keep Out of Reach of Children.



Made in the U.S.A. #34410001

## For the Treatment of Scours and Pneumonia



Net Wt 25 lb (11.34 kg)

## For manufacturing a Type C complete feed to be used in calf milk replacer or starter feed.

#### **Indications for Use**

For calves (up to 250 lbs) feed Super Guard Type B in addition to the calf's regular feed and not as a replacement. Feeding directions are based on individual calves and not based on feeding levels or feeding programs.

### Feeding Directions

One (1) oz of Super Guard Type B medicated feed contains 0.5 gram active ingredient or 5 mg/lb of body weight. To supply 10 mg/lb of body weight for the treatment of bacterial enteritis or bacterial pneumonia add 1 oz per feeding to the calf's regular feed and feed twice per day.

Number of Calves	Amount of Super Guard Type E
1	1 oz
8 16	8 ozs 16 ozs (1 lb)
160	10 lbs

Scoop in each pail holds approximately 1 oz dry weight.

Feed the medicated milk replacer in place of normal calf milk replacer continuously for 7 to 14 days. If symptoms persist after feeding the medicated milk replacer for 2 or 3 days, consult a veterinarian. Continue feeding the medicated milk replacer for 24 to 48 hours after the remission of disease symptoms.

25 lb pail contains 400 feedings.

Manufactured By



A Subsidiary of Merrick Animal Nutrition, Inc. Middleton, WI 53562-0307 USA © 2010

#34410038 Carton: 4 - 3# pails

#34410001 25# Pail #34410002 50# Bag

For treatment of bacterial enteritis (scours) caused by E. coli susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by E. coli susceptible to

For treatment of bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocide* susceptible to oxytetracycline.

#### ACTIVE DRUG INGREDIENTS

Providing the maximum allowable amount of 10 mg/lb of body weight per day, when fed as directed.

#### GUARANTEED ANALYSIS

Crude Protein, minimum 6.0% Crude Fat, minimum 6.0% Crude Fiber, maximum 0.15% Calcium, minimum 0.2%	Vitamin A, minimum 800,000 IU/lb Vitamin D3, minimum 200,000 IU/lb Vitamin E, minimum 800 IU/lb
Calcium, maximum	Keep tightly closed. Store in cool, dry place.

Dried Whey, Kaolin, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Vitamin B12 Supplement, Thiamine Mononitrate, Ascorbic Acid, Biotin, Riboflavin, d-Calcium Pantolhenate, Niacin Supplement, Choline Chloride, Mena-dione Sodium Bisulfite Complex (source of Vitamin K Activity), Folic Acid, Pyridoxine Hydrochloride, Magnesium Sulfate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Cobatt Sulfate, Iron Sulfate, Manganese Sulfate, Copper Sulfate, Cobatt Sulfate, Iron Sulfate, Selaium Idate, Mineral Oil, Glucose, Maltodextrin, Sodium Citrate, Sodium Chloride, Glycine, Potassium Chloride, Calcium Lactate, Ascorbic Acid, Inulin, Monosodium Glutamate, Sodium Aluminosilidate, Dried Bacillus subtilis fermentation product, Dried Lactobacillus acide fermentation product, Dried Sulfobacterium bifidum Fermentation product, Dried Bacillus sulm refermentation product, Dried Bacillus s

#### WARNING

Residue Warning: Withdrawal 5 days before slaughter. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residue. A withdrawal period has not been established for the use in preruminating calves. Do not use in calves to be processed for veal. A discard time has not been established for the use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

For Animal Use Only. Keep Out of Reach of Children.



Made in the U.S.A. #34412001

# For the Treatment of Scours and Pneumonia





Net Wt 25 lb (11.34 kg)

## For manufacturing a Type C complete feed to be used in calf milk replacer or starter feed.

#### Indications for Use

For calves (up to 250 lbs) feed Super Guard Type B in addition to the calf's regular feed and not as a replacement. Feeding directions are based on individual calves and not based on feeding levels or feeding programs.

### Feeding Directions

One (1) oz of Super Guard Type B medicated feed contains 0.5 gram active ingredient or 5 mg/lb of body weight. To supply 10 mg/lb of body weight for the treatment of bacterial enteritis or bacterial pneumonia add 1 oz per feeding to the call's regular feed and feed twice per day.

Number of Calves	Amount of Super Guard Type B
1 8	1 oz 8 ozs
16	16 ozs (1 lb)
160	10 lbs

Scoop in each pail holds approximately 1 oz dry weight.

Feed the medicated milk replacer in place of normal calf milk replacer continuously for 7 to 14 days. If symptoms persist after feeding the medicated milk replacer for 2 or 3 days, consult a veterinarian. Continue feeding the medicated milk replacer for 24 to 48 hours after the remission of disease symptoms.

25 lb pail contains 400 feedings.

Manufactured By



A Subsidiary of Merrick Animal Nutrition, Inc. Middleton, WI 53562-0307 USA © 2010

#34412038 Carton: 4 - 3# pails #34412001 25# Pail #34412002 50# Bag